


Chronic obstructive pulmonary disease

Offer all of the following interventions if appropriate




If the patient still suffering from breathlessness or exercise limitation **offer a SABA as needed.**

The patient is limited by symptoms or suffering exacerbations despite SABA.

Consider if the patient has features suggesting steroid responsiveness:

- ✓ Previous asthma diagnosis or atopy.
- ✓ Substantial variation in FEV1 over time (at least 400ml) or substantial variation in peak flow (at least 20%).
- ✓ A higher blood eosinophil count.



This guideline does not apply to people with COPD and asthma overlap.

[1] Increased risk of pneumonia with inhaled corticosteroid use

Check inhaler technique at every opportunity

Inhaler device is more important than drug choice. Please try to consider device continuity when prescribing different inhalers. **Any inhalers that are on the Pan Mersey formulary can be prescribed.**

NICE COPD Guidelines 2019 suggest ipratropium bromide (SAMA) as an option instead of a SABA

For patients who are using a long acting bronchodilator such as a LABA or LAMA alone whose symptoms are controlled, continue until it is appropriate to change.

Chronic obstructive pulmonary disease

Consider carbon footprint of inhaler choice

Where clinically possible consider use of DPI in preference to MDI

Device	SABA Short Acting Beta 2 Agonist	LAMA+LABA combination Long Acting Muscarinic Antagonist + Long Acting Beta 2 Agonist	ICS + LABA combination Inhaled corticosteroid + Long Acting B2 Agonist	LAMA+LABA/ICS * Long Acting Muscarinic Antagonist + Long Acting Beta 2 Agonist / Inhaled Corticosteroid
DPI Dry Powder Inhaler	Easyhaler® Salbutamol sulphate 100 microgram /dose 1-2 puffs when required	Anoro Ellipta®▼ Umeclidinium 55 microgram / vilanterol 22 microgram 1 puff once daily	Relvar® Ellipta® Fluticasone furoate 92 microgram / vilanterol 22 microgram 1 puff once daily	Trelegy Ellipta®▼ Futicasone furoate 92 microgram / umeclidinium 55 microgram / vilanterol 22 microgram 1 puff once daily
MDI Metered dose inhaler (Spiolto is a soft mist)	Salbutamol CFC free 100 microgram /dose 1-2 puffs when required	Spiolto Respimat®▼ Tiotropium 2.5microgram / olodacterol 2.5 microgram 2 puffs once daily Caution if eGFR < 50	Fostair® Beclometasone 100 microgram / formoterol 6 microgram 2 puffs twice daily Off label if FEV > 50%	Trimbow® Beclometasone 87 microgram / formoterol 5 microgram / glycopyrronium 9 microgram 2 puffs twice daily

See [Pan Mersey Formulary](#) for other first line inhalers

*NB: Offer triple therapy formulations first line but, for individual patients, any appropriate inhalers on the Pan Mersey formulary can be prescribed.